| Barbus lepiney   | <u> </u>                                       |  | Region                    | : 1                       |   |                            |                  |
|--|--|--|---------------------------|---------------------------|---|----------------------------|------------------|
| Taxonomic Authority:   | Pellegrin, 1939                                |  |                           |                           |   |                            |                  |
| Synonyms:  |  |  | Comm                      | on Names:                 |   |                            |                  |
| Order: Cyprinifo   | rmes   |  | Family                    | : Cyprir                  | nidae                                   |                            |                  |
| Notes on taxonomy:   | The taxonomy of barbe group is set in this cou |  | is still uncertain        | and more w                | ork is nee                              | ded before a definitive    | taxonomy of this |
| General Information  | <u>n</u>                                       |  |                           |                           |   |                            |                  |
| Biome  | ☐ Te   | errestrial   | ✓ Freshwate               | er                        | Mar Mar                                 | rine                       |                  |
| <b>Geographic Range of s</b> It is restricted to the Draa Morocco. | •  | o-Saharan zone in  | <b>Habitat</b><br>No data | and Ecolog                | gy Informa                              | ation:                     |                  |
| <b>Conservation Measures</b>                                       | <b>s</b> :                                     |  | Threats                   | s:                        |   |                            |                  |
| None   |  | Climate change is causing drought in all rivers south of the Oum Er Rbia (Azeroual, A. pers comm.) |                           |                           |   |                            |                  |
| Charles namulation info  |  |  | Rbia (A                   | zeroual, A. p             | oers comm                               | 1.)                        |                  |
| Species population info<br>Found to be locally abun                |  | adrio I nere com   | m)                        |                           |   |                            |                  |
| Tourid to be locally abuit   |  |  | 11).                      |                           |   |                            |                  |
|  | Nativ<br>Prese                                 | nce Presence   | Extinct Reir              | ntroduced I               | ntroduced                               | Vagrant                    |                  |
| Country Distribution   |  |  |                           |                           |   |                            |                  |
| Morocco  | <b>✓</b>                                       |  |                           |                           |   |                            |                  |
| Upper Level Habit<br>5.9 Wetlands (inland)                         | tat Preferences<br>- Freshwater Springs an     |  | Score <u>Lowe</u>         | er Level F                | labitat F                               | <u>Preferences</u>         | Score            |
| Major threats  |  |  | Cons                      | ervation                  | Measur                                  | P8                         |                  |
| Code Description of  | throat   | Daet Dresent   | Future Code               |                           |   |                            | In place Needed  |
| 7 Natural disaster   |  |  | <b>✓</b> 3                | Research a                |   | 1103                       |                  |
| 7.1 Drought  | 3  |  | <b>✓</b> 3.1              | Taxonomy                  | actions                                 |                            |                  |
| 7.1 Drought  |  |  | 3.2                       | Population                | numbers a                               | and range                  |                  |
|  |  |  | 3.3                       | Biology and               |   |                            |                  |
|  |  |  | 3.4                       | Habitat sta               | • |                            |                  |
|  |  |  | 3.5                       | Threats                   |   |                            |                  |
|  |  |  | 3.8                       | 3.8 Conservation measures |   |                            |                  |
|  |  |  | 3.9                       | Trends/Mo                 | nitoring                                |                            |                  |
| Utilisation of Species   |  |  |                           |                           |   |                            |                  |
| Purpose/Type of Use  |  | Subsistend   | ce Nationa                | l Internat                |   | Other purpose:<br>lot used |                  |
| Primary forms remove   | d from the wild                                | 100% >75%  | % 51-75%                  | 26-50%                    | <25%                                    | Other forms removed        | from the wild:   |
| Source of specimens i  | n commercial trade                             | 100% >75%  | % 51-75%                  | 26-50%                    | <25%                                    | Other source of speci      | mens:            |
| Trend in wild offtake/h  | arvest in relation to to                       | tal wild population  | on numbers ov             | er last five              | years:                                  |                            |                  |
| Trend in offtake/harve   | st produced through d                          | omestication/cul   | tivation over la          | ast five year             | s:                                      |                            |                  |
| CITES:   |  |  |                           |                           |   |                            |                  |
| Red Listing  |  |  |                           |                           |   |                            |                  |
| Red List Assessment:   | Least Concern (LC)                             |  | F                         | ossibly Exti              | nct                                     |                            |                  |
| Red List Criteria:   |  |  |                           |                           |   |                            |                  |
| Rationale for the Red  | List Assessment: B.                            | lepineyi is widely   | distributed and           | locally abun              | dant (Doa                               | drio, I. pers comm).       |                  |
| <b>Current Population Tre</b>                                      | end: Unknown                                   |  | Date of                   | of Assessm                | <b>ent:</b> 31                          | /10/2004                   |                  |
| Assessor(s): A.J. Cr   | ivelli   |  |                           |                           |   |                            |                  |
| Evaluator: A. Azer   | oual   |  |                           |                           |   |                            |                  |
|  |  |  |                           |                           |   |                            |                  |

## Notes on Red listing: Bibliography

Doadrio, I., Bouhadad, R. & Machordom, A., 1998, Genetic differentiation and biogeography in saharan populations of the genus Barbus (Osteichthyes, Cyprinidae)., Folia Zoologica, , , 47 (Suppl. 1), 7-20, ,

Azeroual, A., 2003, , , Monographie des poissons des eaux continentales du Maroc: systématique, distribution et ecologie., , , , PhD , University Mohammed V, Rabat, Morocco