	<u>tis ohridana</u>	_				Region	: 1					
Taxono	mic Authority: Karaman, 1928	3										
Synonyms: Common Names:												
Order:	Cypriniformes					Family:	Cob	oitidae				
	on taxonomy:						000					
Gener	ral Information											
Biome Terrestrial						✓ Freshwater						
Geographic Range of species: Habitat and Ecology Information:												
								•	he Moraca river where			
FYROM, and the Moraca river and Lake Skadar system in Serbia- Montenegro.						low, and there is a sandy substrate with submerged vegetation. It is also found in the lakeshore.						
Conservation Measures:					Threats:							
None					Gravel extraction, water pollution, habitat destruction							
Species population information:												
Very ab	undant.											
Native - Presence Country Distribution Confirmed			Pres	Native - Presence Extinct Reintroduced Introduced Vagrant Possible								
Albania												
Macedonia, the former Yugoslav Republ				1	_							
	and Montenegro			1	_							
<u> </u>												
	er Level Habitat Preference		(O		core	Lowe	er Level	Habita	t Preferences		Score	
	Vetlands (inland) - Permanent Riv ncludes waterfalls)	ers/Streams	/Creeks	5	1							
•	Vetlands (inland) - Permanent Fre	shwater Lak	es (ove	er 8ha)	1							
Major threats Conservation Measures												
Code				resent						In place	Needed	
1	Habitat Loss/Degradation (huma					3		h actions				
1.3	Extraction	,				3.1	Taxonon	ny				
1.3.6	Groundwater extraction				✓	3.2	Populatio	on numbe	rs and range			
1.3.7	Other		\checkmark	\checkmark	✓	3.3	Biology a	and Ecolog	ду		\checkmark	
1.4	Infrastructure development					3.4	Habitat s	status			\checkmark	
1.4.6	Dams					3.5	Threats				✓	
2	Invasive alien species (directly affecting the species)			\checkmark	✓	3.8						
6	Pollution (affecting habitat and/or species)			\checkmark	3.9 Trends/Monitoring						\checkmark	
6.3				✓	✓							
6.3.1	Agriculture			\checkmark	✓							
6.3.2	Domestic		\checkmark	\checkmark	✓							
6.3.8	Sewage											
9	Intrinsic factors											
9.1	Limited dispersal		✓									
9.9	Restricted range											
	tion of Species											
Purpos	se/Type of Use		Subs	istenc	e	National	Intern	ational	Other purpose: Not used at all.			
Drimar	ry forms removed from the wild	10	0%	>75%	5	1-75%	26-50%	<25%	Other forms remove	d from the wil	d	
	-										а.	
	e of specimens in commercial t		0%	>75%		51-75%	26-50%	<25%	Other source of spe	cimens:		
	in wild offtake/harvest in relation							-				
Trend CITES:	in offtake/harvest produced thr :	ough dome	sticatio	on/cult	ivatio	on over la	ist five ye	ears:				
	L isting I st Assessment : Least Conce	m (I C)				ח 🗌	ossibly E	vtinct				
								AUTOL				
Red List Criteria: Rationale for the Red List Assessment: This species is very abundant throughout its known range with no current signs of decline. It is also											is also	
		known	to be q	uite res	silient	to degrad	ded condi	tions. The	taxonomy and distribut			
0	A Demulation Transfer 11	•	s are co	onfirme	d are		• •	ers comm				
	nt Population Trend: Unknow	/n				Date o	f Assess	inent:	31/10/2004			
Asses		4										
Evalua	ator: Frevhof, J. & Kottelat, M	/1.										

Notes on Red listing:

Bibliography

Economidis, P.S. & Nalbant, T.T., 1996, A study of the loaches of the genera Cobitis and Sabanejewia (Pisces, Cobitidae) of Greece, with description of six new taxa., Trav. Mus. Natl. Hist. Nat. 'Grigore Antipa", , , 36, 295-347, ,

Bohlen, J., Slechtova, V., Sanda, R., Kalous, L., Freyhof, J., Vukic, J. & Mrdak, D., 2003, Cobitis ohridana and Barbatula zetensis in the River Moraça Basin, Montenegro: distribution, habitat, population structure and conservation needs., Folia biologica (Krakow), , , 51 (Suppl.), 147-153,

Kottelat, M., 1997, European freshwater fishes, Biologia, Bratislava, , , 52/ Supplement 5, 1-271, ,