e negudalanidati Dhovinallu

Pnox	inelius pse	udalepidotu	<u>s</u>		Regior	n: 1					
Taxonomic Authority: Bogutskaya & Zupancic, 2003											
Synonyms:					Comm	Common Names:					
Order: Cypriniformes Notes on taxonomy:				Family	Family: Cyprinidae						
	-										
<u>Gener</u> Biome	al Informatio		Terrestrial		Freshwat	or		Marine			
					-	Habitat and Ecology Information:					
Geographic Range of species: It is restricted to a single stream 10km long in the Mostarsko blato (Freyhof, J. pers com), an extensive wetland adjacent to Mostar in Bosnia-Hercegovina. The species is probably spread more widely in the Neretva basin.						It lives in streams with current and clear water.					
Conservation Measures: It is thought that a nature reserve may have been established in this area (Freyhof pers comm).						Threats: No obvious threats (Freyhof pers com).					
•	population info	,									
No data											
	try Distributio	Pro	esence Pre nfirmed Pc	ssible	Extinct Rei	_	_	-			
	and Herzegovina										
		at Preferences		Sco.		er Level	Habita	<u>t Preferences</u>		Score	
	/etlands (inland) - ncludes waterfalls	Permanent Rivers/ ;)	Streams/Cree	ks	1						
Major	Major threats				Con	Conservation Measures					
Code	Description of	threat	Past	Present Fu	ture Code	Conserv	ation me	asures	In place	Needed	
1	Habitat Loss/De	gradation (human in	duced)		✔ 1	Policy-ba	ased actio	ns		\checkmark	
1.3	Extraction				✔ 3	Researc	h actions		\checkmark	\checkmark	
1.3.6	Groundwater ex	traction			✔ 3.1	Taxonom	ıy		\checkmark		
2		pecies (directly affect	ting the		✓ 3.2	•		rs and range			
7	species) Natural disaster	8			3.3 ✓ 3.4		and Ecolo	ду			
, 7.1	Drought	5				Habitat s	tatus				
9	Intrinsic factors					Threats					
9.1	Limited dispersa	I					ation mea				
9.10	Other	-					Aonitoring				
					▲ 4 4.1		ance/Cons	ased actions			
					4.1	Protected		Servation			
					4.4.2					\checkmark	
					4.4.3						
Utilisat	tion of Species										
Purpos	se/Type of Use		Sub	sistence	Nationa	al Intern	ational	Other purpose: Not used at all.			
Primary forms removed from the wild			100%	>75%	51-75%	26-50%	<25%	Other forms remov	ed from the w	ild:	
Source of specimens in commercial trade			9 100%	>75%	51-75%	26-50%	<25%	Other source of sp	ecimens:		
Trend	in wild offtake/h	arvest in relation to	total wild po	opulation	numbers o	ver last fiv	e years:				
Trend		t produced throug	h domesticat	ion/cultiva	ation over l	ast five ye	ars:				
Red L	isting										
Red List Assessment: Vulnerable (VU)				Possibly Ex	xtinct						
	st Criteria: D2										
Rationale for the Red List Assessment: The species appears to have r known from only one location, an extensive wetland adjacent more widely in the Neretva bas						km stretch	of stream	n (Freyhof, J. pers com	n) in Mastarsko	blato,	
Curren	t Population Tre	nd: Unknown	more widely			of Assess	ment [.]	31/10/2004			
Assess	•				Duto						

Evaluator:	J. Freyhof
------------	------------

Notes on Red listing:

Bibliography

Bogutskaya, N.G. & Zupancic, P., 2003, Phoxinellus pseudalepidotus (Teleostei: Cyprinidae), a new species from the Neretva basin with an overview of the morphology of Phoxinellus species of Croatia and Bosnia-Herzegovina., Ichthyol. Explor. Freshwaters, , , 14, 369-383, ,