Chalcides cha	<u>alcides</u>			Region: 7						
Taxonomic Authority	y: (Linnaeus, 1758)									
Synonyms:		Common Names:								
Chalcides mertensi	Klausewitz,	1954		Three-toed Skin	k	English				
					ed Skink	English	English			
				luscengola		Italian				
Order: Sauria	a			Family: So	incidae					
Notes on taxonomy:			si as a s	separate species	, rathan than a	as a subspecies of	Chalcides chalcides,			
	following Caputo (1993).								
General Informa	<u>tion</u>									
Biome	✓ Ter	rrestrial	Fre	eshwater	☐ Mar	ine				
Geographic Range of	of species:			Habitat and Ec	ology Informa	ition:				
the Po Valley), include on a number of smaller	s occurs in Italy (where it is ng on the islands of Sicily, S er islands. In North Africa it i northern Tunisia and north el up to 1,270m asl.	Sardinia and Elba is found in	and can	include, grassla cork-oak forest	nd, meadows, and the edges aded areas. Tl	areas close to stre of cultivated areas	vegetation. Habitats eams, hedges, open s. It can also be found rth to three to thirteen			
Conservation Measu	ires:			Threats:						
protected areas.	on the Bern Convention. It c	occurs in a numbe		but habitat loss threat to the spe the freshwater/o	through intens cies in some a oastal zones o	ification of agricult				
Species population It is common in suitab										
	Native Preser		Cutio	at Daintraduae	d Introduced	Vagrant				
Country Distribu			⊏XUII	ct Reintroduce	u miloduced	Vagrant				
Algeria	<u>∡tion</u>									
Italy	▼			_						
Libyan Arab Jamahiri										
Tunisia	,			_						
FAO Marine Hab	Nativ Prese	nce Presence	Extin	nct Reintroduce	d Introduced					
Major Lakes										
Major Rivers										
	oitat Preferences	S		Lower Leve	l Habitat P	<u>references</u>	Score			
1.4 Forest - Temper			1							
3.4 Shrubland - Ter	•	4-4:	1							
	diterranean-type Shrubby Ve	egetation	1							
4.4 Grassland - Ter5.2 Wetlands (inlan Rivers/Streams/	d) - Seasonal/Intermittent/Irr	regular	1							
11.2 Artificial/Terrest			1							
11.4 Artificial/Terrest	rial - Rural Gardens		1							
Major threats				Conservation	n Measure	es				
Code Description	of threat	Past Present		Code Conserv			In place Needed			
2000	Degradation (human induce		✓		ased actions		pides 1166des			
1.1 Agriculture			✓	1.2 Legislat						
1.1.1 Crops			✓	1.2.1 Develop						
1.1.1.3 Agro-industry	farming		✓	1.2.1.1 Internati						
•	development		✓	1.2.1.2 Nationa						
1.4.2 Human settle		<u> </u>	✓	1.2.2 Impleme	entation		V			

1.4.3 6 6.2 6.2.1	Tourism/recreation Pollution (affecting habitat and/or special pollution Agriculture	ecies)		1.2 3 3.3 3.5 3.5 4 4.6 4.6 4.6 4.6	Taxonor Threats Trends/I Habitat a Mainten: Protecte 2 Establisl	h actions ny Monitoring and site-bance/Cons d areas nment	I ased actions						
Utilisa	tion of Species												
Purpose/Type of Use		Subsistence		. Nati	National International		Other purpose:						
Primary forms removed from the wild		100% >75% 51-75% 26-50% <25% Other forms removed from th						rom the wild	l:				
Source of specimens in commercial trade		100%	>75%	51-75	26-50%	26-50% <25% Other source of specia							
Trend in wild offtake/harvest in relation to total wild population numbers over last five years: Trend in offtake/harvest produced through domestication/cultivation over last five years: CITES:													
Red Listing Red List Assessment: Least Concern (LC)													
Rationale for the Red List Assessment: Listed as Least Concern in view of its wide distribution, tolerance of a degree of habitat modification, presumed large population, and because it is unlikely to be declining fast enough to qualify for listing in a more threatened category.													
Currer	nt Population Trend: Decreasing	Date of Assessment: 12/17/2004											
Assessor(s): Jose Antonio Mateo Miras, Marc Cheylan, M. Saïd Nouira, Ulrich Joger, Paulo Sá-Sousa, Valentin Pérez-Mellado													
Notes	on Red listing:												

Bibliography

Arnold, E.N., 2003, , , Reptiles and amphibians of Europe., , , 288 pp., Princeton University Press., Princeton and Oxford.

Gasc, J.-P., Cabela, A., Crnobrnja-Isailovic, J., Dolmen, D., Grossenbacher, K., Haffner, P., Lescure, J., Martens, H., Martínez-Rica, J.P., Maurin, H., Oliveira, M.E., Sofianidou, T.S., Veith, M. and Zuiderwijk, A., 1997, , , Atlas of Amphibians and Reptiles in Europe., , , pp. 494, Societas Europaea Herpetologica and Musée National d'Histoire Naturelle, Paris

Caputo, V. and Odierna, G., 1992, Karyological differentiation between two forms of the Chalcides chalcides complex (Scincidae)., Amphibia-Reptilia, , , 13, 193-196, ,

Busack, S.D., 1986, Taxonomic implications of biochemical and morphological differentiationin Spanish and Moroccan populations of three-toed skinks, Chalcides chalcides (Lacertilia, Scincidae)., Herpetologica, , , 42(2), 230-236, ,

Schleich, H.H., Kästle, W. and Kabisch, K., 1996, , , Amphibians and Reptiles of North Africa. , , 627 pp., Koeltz., Koenigstein.

Bons, J. and Geniez, P., 1996, , , Amphibiens et Reptiles du Maroc (Sahara Occidental compris), Atlas biogéographique., , , 320 pp., Asoc. Herpetol. Espanola., Barcelona

Caputo, V., 1993, Taxonomy and evolution of the Chalcides chalcides complex (Reptilia, Scincidae) with description of two new species., Museo Regionale de Scienze Naturali Bollettino (Turin), , , 11(1):, 47-120, ,