Chalcides montanus Taxonomic Authority: Werner, 1931			Re	gion: 7			
Synonyms:				Common Names: Mountain Skink English			
Order: Sauria			Fa	mily: So	incidae	-	
Notes on taxonomy:							
General Information Biome	✓ Terrestri	al	Fresh	nwater	N	Marine	
Geographic Range of species: This species is endemic to the High Atlas found between 2,300 and 2,830m asl.	Mountains o	f Morocco. I	t is Th bu	shland, fields	found in coo , meadows	rmation: ol montane cedar planta and damp areas close to s a viviparous species.	
Conservation Measures: It has been recorded from the Toubkral N	ational Park.		Alt ha	bitat destructi	on through	believed to be significan removal of firewood is the ts of its range.	
Species population information: It is a rare species that is not thought to b range.	e abundant v	vithin its narı	row				
Country Distribution	Native - Presence Confirmed	Native - Presence Possible	Extinct	Reintroduce	d Introduc	ed Vagrant	
Morocco	✓						
FAO Marine Habitats	Native - Presence Confirmed	Native - Presence Possible	Extinct	Reintroduce	ed Introduc	ced	
Major Lakes Major Rivers							
Upper Level Habitat Preference	<u>es</u>	S		ower Leve	l Habitat	Preferences	Score
1.4 Forest - Temperate3.4 Shrubland - Temperate			1 1				
6 Rocky areas (eg. inland cliffs, mour	itain peaks)		1				
11.2 Artificial/Terrestrial - Pastureland			1				
Major threats			С	onservatio	n Measu	ıres	
CodeDescription of threat1Habitat Loss/Degradation (human1.3Extraction1.3.3Wood1.3.3.1Small-scale subsistence1.4Infrastructure development1.4.3Tourism/recreation	n induced)	ast Present V V V V V V V V V V V V V V V V V V	Future	Taxonor Populati Biology Habitat	ch actions my ion numbers and Ecolog	s and range	In place Needed
Utilisation of Species							
Purpose/Type of Use	;	Subsistenc	e Nati	onal Interr	national	Other purpose:	
Primary forms removed from the wild	100%	% >75%	51-75	% 26-50%	<25%	Other forms removed	from the wild:
Source of specimens in commercial tra	ade 100%	% >75%	51-75	% 26-50%	<25%	Other source of speci	mens:
Trend in wild offtake/harvest in relation Trend in offtake/harvest produced thro CITES:					-		

Red List Criteria:

Rationale for the Red List Assessment:

Listed as Near Threatened because, although its Extent of Occurrence is less than 20,000 km2, it

occurs in a habitat that is not under significant threat, and so it is probably not in decline.

Current Population Trend: Stable

Date of Assessment:

12/17/2004

Assessor(s): Jose Antonio Mateo Miras, Ulrich Joger, Tahar Slimani, El Hassan El Mouden, Philippe Geniez

Notes on Red listing:

Bibliography

Bons, J. and Geniez, P., 1996, , , Amphibiens et Reptiles du Maroc (Sahara Occidental compris), Atlas biogéographique., , , 320 pp., Asoc. Herpetol. Espanola., Barcelona

Caputo, V. and Pêrez-Mellado, J., 1992, A new species of Chalcides (Reptilia: Scincidae) from northeastern Morocco., Bollettino de Zoologia, , , 59(3):, 335-342, ,

Schleich, H.H., Kästle, W. and Kabisch, K., 1996, , , Amphibians and Reptiles of North Africa., , , 627 pp., Koeltz., Koenigstein.

Mateo, J.A., Geniez, P. and Bons, J., 1995, Saurians of the genus Chalcides Laurenti 1768 (Reptilia, Scincidae) in Morocco, I. review and distribution., Rev. Española Herp., , , 9, 7-36, ,